



Korean American Community Services

PFIZER-BIONTECH COVID-19 VACCINE

MODERNA COVID-19 VACCINE

JANSSEN COVID-19 VACCINE (ALSO KNOWN AS JOHNSON & JOHNSON)

AUTHORIZATION

State of California ☒
U.S. Food and Drug
Administration (FDA) ☒

State of California ☒
U.S. Food and Drug
Administration (FDA) ☒

State of California ☒
U.S. Food and Drug
Administration (FDA) ☒

VACCINE TYPE/VACCINE DELIVERY METHOD

mRNA

mRNA

Conventional
Viral Vector

NUMBER OF DOSES

2

2

1

EFFECTIVENESS

95% effective at preventing
COVID-19 illness after
second dose

94.1 effective at preventing
COVID19 illness after
second dose

74% effectiveness against
moderate and severe cases of
COVID-19, with levels rising to
86% effectiveness in preventing
severe cases after 4 weeks

MOST COMMON SIDE EFFECTS

In the arm where you got
the shot: pain swelling,
redness

Throughout the rest of your
body: chills, tiredness,
headache

In the arm where you got
the shot: pain swelling,
redness

Throughout the rest of your
body: chills, tiredness,
headache

In the arm where you got
the shot: pain swelling,
redness

Throughout the rest of
your body: chills,
tiredness, headache

SECOND DOSE DUE

21 days
after first dose

28 days
after first dose

Second dose not
needed